



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Changing glucocorticoid action

**Citation for published version:**

Chapman, KE, Coutinho, AE, Zhang, Z, Kipari, T, Savill, JS & Seckl, JR 2013, 'Changing glucocorticoid action: 11-hydroxysteroid dehydrogenase type 1 in acute and chronic inflammation', *Journal of Steroid Biochemistry and Molecular Biology*, vol. 137, pp. 82-92. <https://doi.org/10.1016/j.jsbmb.2013.02.002>

**Digital Object Identifier (DOI):**

[10.1016/j.jsbmb.2013.02.002](https://doi.org/10.1016/j.jsbmb.2013.02.002)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Journal of Steroid Biochemistry and Molecular Biology

**Publisher Rights Statement:**

Gold Open Access paid so please attach the publisher final pdf

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Accepted Manuscript

Title: Changing glucocorticoid action: 11 $\beta$ -hydroxysteroid dehydrogenase type 1 in acute and chronic inflammation

Authors: Karen E. Chapman, Agnes E. Coutinho, Zhenguang Zhang, Tiina Kipari, John S. Savill, Jonathan R. Seckl



PII: S0960-0760(13)00021-6  
DOI: <http://dx.doi.org/doi:10.1016/j.jsbmb.2013.02.002>  
Reference: SBMB 3940

To appear in: *Journal of Steroid Biochemistry & Molecular Biology*

Received date: 26-9-2012  
Revised date: 22-1-2013  
Accepted date: 4-2-2013

Please cite this article as: K.E. Chapman, A.E. Coutinho, Z. Zhang, T. Kipari, J.S. Savill, J.R. Seckl, Changing glucocorticoid action: 11 $\beta$ -hydroxysteroid dehydrogenase type 1 in acute and chronic inflammation, *Journal of Steroid Biochemistry and Molecular Biology* (2013), <http://dx.doi.org/10.1016/j.jsbmb.2013.02.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Title: Changing glucocorticoid action: 11 $\beta$ -hydroxysteroid dehydrogenase type 1 in acute and chronic inflammation.**

**Authors: Karen E Chapman<sup>1</sup>, Agnes E Coutinho<sup>1</sup>, Zhenguang Zhang<sup>1</sup>, Tiina Kipari<sup>1</sup>, John S Savill<sup>2</sup>, Jonathan R Seckl<sup>1</sup>**

**Affiliations:** <sup>1</sup>University/BHF Centre for Cardiovascular Sciences and <sup>2</sup>MRC Centre for Inflammation Research, The Queen's Medical Research Institute, University of Edinburgh, 47 Little France Crescent, Edinburgh, EH16 4TJ, UK.

**Correspondence:** Karen E Chapman.  
University/BHF Centre for Cardiovascular Sciences, The Queen's Medical Research Institute, University of Edinburgh, 47 Little France Crescent, Edinburgh, EH16 4TJ, UK  
Tel: 44-131-242-6736; FAX: 44-131-242-6779  
e-mail: [Karen.Chapman@ed.ac.uk](mailto:Karen.Chapman@ed.ac.uk)

**Disclosures:** Jonathan R Seckl holds IP on the use of 11 $\beta$ -HSD1 inhibitors in diabetes, atherosclerotic disease and age-associated cognitive impairment. None of the other authors have anything to disclose.

**Abbreviations:** 11 $\beta$ -HSD, 11 $\beta$ -hydroxysteroid dehydrogenase; H6PD, hexose-6-phosphate dehydrogenase; TNF- $\alpha$ , tumour necrosis factor- $\alpha$ ; LPS, lipopolysaccharide; IL-, interleukin; C/EBP, CCAAT/enhancer binding protein; NF- $\kappa$ B, nuclear factor kappa-light-chain-enhancer of activated B cells; EGR-1, early growth response-1; HPA, hypothalamic-pituitary-adrenal; MCP, monocyte chemotactic protein; VCAM, vascular cell adhesion molecule.

**Keywords:** glucocorticoid, mineralocorticoid, 11 $\beta$ -hydroxysteroid dehydrogenase, macrophage, inflammation, arthritis

## ABSTRACT

Since the discovery of cortisone in the 1940s and its early success in treatment of rheumatoid arthritis, glucocorticoids have remained the mainstay of anti-inflammatory therapies. However, cortisone itself is intrinsically inert. To be effective, it requires conversion to cortisol, the active glucocorticoid, by the enzyme 11 $\beta$ -hydroxysteroid dehydrogenase type 1 (11 $\beta$ -HSD1). Despite the identification of 11 $\beta$ -HSD in liver in 1953 (which we now know to be 11 $\beta$ -HSD1), its physiological role has been little explored until recently. Over the past decade, however, it has become apparent that 11 $\beta$ -HSD1 plays an important role in shaping endogenous glucocorticoid action. Acute inflammation is more severe with 11 $\beta$ -HSD1-deficiency or inhibition, yet in some inflammatory settings such as obesity or diabetes, 11 $\beta$ -HSD1-deficiency/inhibition is beneficial, reducing inflammation. Current evidence suggests both beneficial and detrimental effects may result from 11 $\beta$ -HSD1 inhibition in chronic inflammatory disease. Here we review recent evidence pertaining to the role of 11 $\beta$ -HSD1 in inflammation.

## Introduction

The discovery of the anti-inflammatory effects of cortisone, a glucocorticoid hormone, by Hench and colleagues in the 1940s, opened the door to the longest and most successful drug development programme in history. Glucocorticoids remain the most widely prescribed treatment for inflammatory disease. They potently affect both immune and non-immune cells, shaping their responses. Glucocorticoid actions are highly dependent on context and can be very different during acute and chronic inflammation. In the short term at least, many of their effects promote the resolution of inflammation. Several years ago, we hypothesised that the glucocorticoid metabolising enzyme, 11 $\beta$ -hydroxysteroid dehydrogenase type 1 (11 $\beta$ -HSD1), is induced early during an inflammatory response and shapes its subsequent trajectory [1]. How well has that hypothesis stood the test of time? Reasonably well as it turns out, but not in quite the way we had envisaged.

## Glucocorticoids and inflammation

Synthetic glucocorticoids exert potent anti-inflammatory and immunosuppressive effects and are widely prescribed to treat both acute and chronic inflammation. Yet the well known side effects of glucocorticoid excess include type 2 diabetes, visceral obesity, hypertension and atherosclerosis which are themselves, somewhat paradoxically, inflammatory conditions. Quite how glucocorticoids provoke inflammatory metabolic diseases at the same time as suppressing chronic inflammatory conditions such as rheumatoid arthritis or inflammatory bowel disease remains unclear. It is likely to involve more complex mechanisms than the commonly held view that the “adverse” metabolic effects involve gene activation by glucocorticoid receptor (GR), whereas the “beneficial” anti-inflammatory effects rely on gene repression. Fully understanding how glucocorticoids cause “metabolic inflammation” will be crucial for the development and optimal exploitation of future anti-inflammatory therapies, which could manipulate glucocorticoid action in a more sophisticated manner than current therapies.

Understanding the role of endogenous glucocorticoids during inflammation is key to achieving this aim. Endogenous glucocorticoids are vital to survive trauma or certain bacterial infections; they suppress pro-inflammatory cytokine production, binding to GR in immune cells to prevent potentially lethal overshoot of immune responses [2, 3]. Acutely, circulating pro-inflammatory cytokines are a potent stimulus to the hypothalamic-pituitary-adrenal (HPA) axis to increase endogenous glucocorticoid production [4, 5]. However, this normal response is lost or attenuated in chronic inflammation [6]. In this respect, the treatment of chronic inflammatory disease with exogenous glucocorticoids can be regarded as replacement therapy for an inadequate endogenous glucocorticoid response [7].

Acute inflammation is an immediate response of the body to injury or infection that serves to remove the injurious stimulus, then restore homeostasis by removal of dead and damaged cells/tissues and engagement of repair processes. It is initiated at the site of injury by the release of proinflammatory mediators such as bioactive amines, lipids and cytokines: typically tumour necrosis factor (TNF)- $\alpha$  and interleukin (IL)-1. These cause vasodilation, increase vascular permeability allowing exudation of plasma, and elicit leukocyte recruitment, activation and emigration from the microcirculation to the damaged tissue. The initial response is typically predominated by neutrophils, which are replaced by monocytes/macrophages during the resolution and repair stages. Resolution of acute inflammation requires the engagement of mechanisms early in the inflammatory response that shape the subsequent resolution (reviewed, [8-10]). Chronic

inflammation results from persistence of the initiating stimulus with associated lymphocyte and macrophage activation. Excessive tissue damage contributes to continuing inflammation, failure of resolution and dysregulated repair processes such as angiogenesis and fibrosis and can thus form a “vicious” cycle. Whilst acute inflammation frequently occurs and is contained entirely at the local level, chronic inflammation invariably involves a systemic response.

Glucocorticoids limit acute inflammation. They repress a large number of proinflammatory genes, including pro-inflammatory cytokines and chemokines, cell adhesion molecules and enzymes involved in the initiation and/or maintenance of inflammation, many of which are over-expressed in chronic non-resolving inflammation. Conversely, they activate a number of genes encoding anti-inflammatory mediators, such as IL-10 and annexin I (reviewed, [11-13]). Thus, acutely, glucocorticoids inhibit the initial vasodilation and increased vascular permeability during inflammation. They also alter the balance between survival and apoptosis of leukocytes as well as their distribution between the circulation and immune tissues and they decrease leukocyte emigration into sites of injury [13-18]. Importantly, glucocorticoids potently influence the differentiation and phenotype of immune cells, especially monocytes/macrophages and T lymphocytes, thereby polarising, or shaping, immune responses [19]. Glucocorticoid treatment of human monocytes promotes an anti-inflammatory, pro-resolution phenotype, characterised by high migratory and phagocytic capacity, expression of CD163 (haemoglobin scavenger receptor) and high production of IL-10 [20-23]. Similarly, in mice, pro-resolving macrophage functions are enhanced by glucocorticoid treatment [24, 25], thus shaping the trajectory of an inflammatory response and its outcome. Because glucocorticoids inhibit production of “Th1” cytokines, which promote a cell-mediated immune response (activation of phagocytes, antigen-specific T lymphocytes) whilst preserving or promoting “Th2” cytokine production (aiding antibody production), they also shape the adaptive immune response.

Most research on the anti-inflammatory actions of glucocorticoids has utilised dexamethasone, a potent synthetic glucocorticoid with powerful immunosuppressive properties. However, the endogenous glucocorticoids, cortisol (the main glucocorticoid in humans) and corticosterone (in rats and mice), are immunomodulatory rather than immunosuppressive [14, 26], particularly when administered at physiologically relevant concentrations. Indeed, low doses of corticosterone stimulate whereas higher doses suppress macrophage activity [27]. This could, in part, reflect the higher affinity binding of endogenous glucocorticoids to the mineralocorticoid receptor (MR) (dexamethasone poorly activates MR [28]) than to GR as both are expressed in macrophages [27, 29, 30]. However, whereas knock-down or antagonism of GR in macrophages abrogates responses to both high and low doses of corticosterone, knock-down or antagonism of MR has little effect [27], suggesting GR-mediated effects, at least in the rat macrophages tested. The interplay between GR and MR in macrophage function and polarisation is likely to be complex (see below).

#### *11 $\beta$ -hydroxysteroid dehydrogenases modulate glucocorticoid action*

Endogenous glucocorticoids differ from dexamethasone in another important respect; dexamethasone is not inactivated by 11 $\beta$ -HSD activity [31] whereas endogenous glucocorticoids are substrates for the 11 $\beta$ -HSDs, which are important modulators of physiological glucocorticoid action [32]. The 11 $\beta$ -HSD “shuttle” interconverts active glucocorticoids (cortisol, corticosterone) with their 11-keto forms (cortisone, 11-dehydrocorticosterone), which bind poorly to receptors and are therefore intrinsically inert. In intact cells, 11 $\beta$ -HSD1 exhibits oxo-reductase activity, converting cortisone

and 11-dehydrocorticosterone into active cortisol and corticosterone respectively, increasing intracellular glucocorticoid levels. In contrast, 11 $\beta$ -HSD2 is exclusively a dehydrogenase, inactivating cortisol and corticosterone. Expression of 11 $\beta$ -HSD2 is largely restricted to mineralocorticoid-target tissues, most notably the distal nephron of the kidney where it protects the non-selective MR from activation by glucocorticoids, conferring aldosterone-specificity upon MR, which is otherwise a high affinity glucocorticoid receptor [33, 34]. Of the synthetic glucocorticoids in widespread use as anti-inflammatory drugs, it is worth noting that some, including prednisone/prednisolone, are excellent substrates for the 11 $\beta$ -HSDs.

11 $\beta$ -HSD1 is widely expressed, including in immune cells, where its activity is dynamically regulated depending on cell activation state (reviewed, [35]). 11 $\beta$ -HSD1 is up-regulated upon activation of monocytes/macrophages, neutrophils or lymphocytes [35, 36] (and see Figure 1). Circulating leukocytes in mice and healthy humans do not express 11 $\beta$ -HSD2 [1, 37]. Both 11 $\beta$ -HSD isozymes are regulated by pro-inflammatory signalling in non-immune cells (see below for details).

#### *11 $\beta$ -HSD1 expression in monocytes/macrophages depends on cell activation state*

Monocytes and macrophages are essential during an inflammatory response. In response to diverse environmental signals, “resting” or naïve macrophages adopt distinct phenotypes. These are broadly categorised based on *in vitro* experiments into two states, M1 (or classically activated) and M2 (or alternatively activated) (reviewed [38, 39]). M1 macrophages, induced by interferon- $\gamma$  and Toll-like receptor (TLR) activation (eg by lipopolysaccharide, LPS), are vital for host defence, expressing pro-inflammatory cytokines, inducible nitric oxide synthase (iNOS) and demonstrating strong microbicidal activity. M2 macrophages, polarised with IL-4 and/or IL-13, restore homeostasis in the repair phase of inflammation. They are also vital for parasite elimination. Other stimuli induce M2-like anti-inflammatory phenotypes, distinct from IL-4/IL-13 polarised macrophages. Macrophage phenotype *in vivo* may be more complex and heterogeneous [40], especially macrophages with M2-like characteristics, reflecting the diversity of signalling and context *in vivo*. Glucocorticoids restrain M1 macrophages, dampening pro-inflammatory cytokine expression, and in naïve monocytes/macrophages, induce a highly phagocytic, highly motile, M2-like phenotype [21, 24, 41]. Conditional deletion of GR in macrophages increases pro-inflammatory cytokine production and mortality following LPS administration [3, 42]. Conversely, conditional deletion of MR in macrophages promotes polarisation to an alternatively activated (M2) phenotype [43], suggesting a possible reciprocal relationship between GR and MR activation in macrophages. There is therefore considerable potential for 11 $\beta$ -HSD1 (which can potentially supply ligand to either receptor) to modulate monocyte/macrophage phenotype by increasing intracellular glucocorticoid levels, even in the absence of elevated circulating glucocorticoid levels.

Expression of 11 $\beta$ -HSD1 is low in circulating mouse leukocytes but is higher in macrophages [44]. Though negligible in non-stimulated human monocytes, 11 $\beta$ -HSD1 expression is induced upon differentiation into resting or naïve (ie unstimulated) macrophages [37]. M1 polarisation of naïve macrophages with LPS further induces 11 $\beta$ -HSD1 (Figure 1). In contrast, polarisation to an M2 phenotype with IL-4 has little effect on 11 $\beta$ -HSD1 expression [45, 46]. However, in human monocytes differentiated into macrophages in the presence of IL-4 (which may induce a distinct anti-inflammatory macrophage phenotype from M2 polarisation of resting macrophages), 11 $\beta$ -HSD1 activity is as high or higher than in M1, and is further increased by peroxisome proliferator-activated receptor (PPAR)- $\gamma$  activation [47]. In contrast, in mouse bone marrow-derived macrophages (resting macrophages) PPAR $\gamma$  agonists

down-regulate 11 $\beta$ -HSD1 expression [47]; whether this reflects a mouse/human species difference or the different macrophage phenotypes (resting mouse macrophages *versus* human macrophages differentiated in the presence of IL-4) is currently unclear. Nevertheless these studies illustrate a complex dependence of 11 $\beta$ -HSD1 expression upon macrophage activation state. The significance is currently unknown but might reflect (or influence) differences in energy metabolism between glycolytic M1 and oxidative M2 macrophages [48, 49]. Recent evidence suggests manipulation of glucose metabolism in macrophages directly alters polarisation [49]. Whether alterations in 11 $\beta$ -HSD1 expression influence macrophage glucose metabolism, for example through the coupling of 11 $\beta$ -HSD1 oxo-reductase activity to hexose 6-phosphate activity in the endoplasmic reticulum (see below) is an important question to address as it may directly affect polarisation or the extent of activation of macrophages. Dynamic regulation of 11 $\beta$ -HSD1 in macrophages could therefore be crucial to the ability to shape an ongoing inflammatory response, either through intracellular regeneration of glucocorticoids or indirectly by diversion of glucose-6-phosphate (Figure 2). Evidence for dynamic regulation of 11 $\beta$ -HSD1 during an inflammatory response *in vivo* comes from the rapid induction of 11 $\beta$ -HSD1 activity in neutrophils and monocytes/macrophages during sterile peritonitis in mice; 11 $\beta$ -HSD1 activity decreases as the inflammation resolves [1, 36]. The latter is possibly an active process; 11 $\beta$ -HSD1 activity is rapidly down-regulated in macrophages that have phagocytosed apoptotic neutrophils [35], a highly pro-resolution process [50]. This reasoning led to the hypothesis that the early induction of 11 $\beta$ -HSD1 in macrophages increases glucocorticoid action within these cells, promoting an anti-inflammatory phenotype and leading to more rapid resolution of inflammation [1, 51].

#### 11 $\beta$ -HSD1 in acute inflammation; regulation

In most animal models of acute inflammation 11 $\beta$ -HSD1 activity is up-regulated in the inflamed tissue, whereas 11 $\beta$ -HSD2 (if expressed at all) is down-regulated. This is true of the inflamed colon and the arthritic joint [52-54], but not the vasculature [55]. This switch in the balance of 11 $\beta$ -HSD1 and 2 activities is predicted to increase paracrine/autocrine glucocorticoid action, though this has not been directly tested. Induction of 11 $\beta$ -HSD1 (and repression of 11 $\beta$ -HSD2) at inflamed sites is probably due to local release of the pro-inflammatory cytokines IL-1 and TNF- $\alpha$  which stimulate transcription of the 11 $\beta$ -HSD1 gene promoter through increased binding of the transcriptional regulators CCAAT/enhancer binding protein (C/EBP) $\alpha$  and nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B) [56-58] and repress the 11 $\beta$ -HSD2 gene promoter through an early growth response (EGR)-1 and NF- $\kappa$ B-dependent mechanism [59]. Normally glucocorticoids antagonise TNF- $\alpha$  or IL-1 action, but they act together with the pro-inflammatory cytokines to synergistically increase 11 $\beta$ -HSD1 expression in a variety of cell types [60-63]. This is predicted to amplify the effect of glucocorticoid within a given cell or tissue, more rapidly promoting the repair and resolution phase. Whether 11 $\beta$ -HSD1 expression in inflammatory cells is regulated by similar mechanisms is an interesting question. Neither TNF- $\alpha$  nor IL-1 $\alpha$  affect 11 $\beta$ -HSD1 activity in monocytes [37] and the signalling pathways that regulate macrophage 11 $\beta$ -HSD1 expression have not been characterised. C/EBP $\alpha$ , a key regulator of 11 $\beta$ -HSD1 transcription in a variety of cell types [56, 57, 64-68], mediates M2 polarisation and arginase expression [69] yet also plays a role in pro-inflammatory cytokine expression in M1 macrophages [70]. However, genetic deletion of C/EBP $\alpha$  abolishes both the liver-enriched inhibitor protein (LIP) and liver-enriched activator protein (LAP) C/EBP $\alpha$  isoforms, the balance of which potentially influences 11 $\beta$ -HSD1 mRNA levels *in vivo* [71] and also regulates osteoclast differentiation [72], a process akin to macrophage differentiation. The C/EBP $\alpha$ -LIP:LAP ratio is regulated by mTOR [73], an integrator of cellular nutrient

and energy metabolism, that is downstream of phosphatidylinositol 3-kinase (PI3K) and Akt, both capable of polarising macrophages [74, 75]. Plausibly, the C/EBP $\beta$ -LIP:LAP ratio differs according to the activating stimulus and may govern the expression level of 11 $\beta$ -HSD1 in polarised macrophages. The coupling within the endoplasmic reticulum of 11 $\beta$ -HSD1 activity to the supply of NADP(H) cofactor generated by hexose-6-phosphate dehydrogenase (H6PD) [76-78] is particularly intriguing in this respect, as it raises the possibility that cellular glucose availability and flux through the endoplasmic reticulum pentose phosphate pathway (the first 2 steps of which are catalysed by H6PD) controls 11 $\beta$ -HSD1 activity [79] which may therefore differ irrespective of expression levels in M1 and M2 macrophages.

#### 11 $\beta$ -HSD1 in acute inflammation; function

Based on the expression of 11 $\beta$ -HSD1 in macrophages, its induction early during an inflammatory response and the well-known anti-inflammatory effects of glucocorticoids, it was anticipated that 11 $\beta$ -HSD1 deficiency or inhibition would attenuate local glucocorticoid production and thus worsen acute inflammation. This is indeed what is seen in 11 $\beta$ -HSD1-deficient (*Hsd11b1*<sup>-/-</sup>) mice, with more severe LPS-induced endotoxaemia (classically repressed by glucocorticoids [3, 80]), an earlier onset of inflammation in the K/BxN serum transfer model of inflammatory arthritis and more inflammatory cells (both neutrophils and monocytes/macrophages) recruited in sterile peritonitis or pleuritis and in the injured myocardium following myocardial infarction [81-83] (and see Figure 2). This increase in inflammation could reflect greater recruitment and/or delayed clearance/apoptosis of neutrophils [1, 84]. In support of the latter, *Hsd11b1*<sup>-/-</sup> mice show delayed macrophage acquisition of phagocytic capacity for apoptotic neutrophils as well as an increase in the number of free apoptotic neutrophils during sterile peritonitis, although surprisingly the peritonitis resolves at the same time as in wild-type mice [1]. Also surprising was the finding that despite the increased inflammation early following myocardial infarction or possibly because of it, heart function post-infarction is much better preserved in *Hsd11b1*<sup>-/-</sup> mice than in controls. Underlying the improved recovery from myocardial infarction is an increased angiogenic response to injury [85], probably as a consequence of an earlier accumulation of reparative M2 (Ym1<sup>+</sup>) macrophages and higher levels of the pro-angiogenic cytokine IL-8 in the hearts of *Hsd11b1*<sup>-/-</sup> mice [82]. It will be important to determine how generally this accelerated switch in macrophage phenotype from M1 to M2 applies to inflammation in *Hsd11b1*<sup>-/-</sup> mice; so far it has only been reported in myocardial infarction and M2-like polarisation is not a general feature of 11 $\beta$ -HSD1-deficient macrophages, at least *in vitro* [1, 81] or *in vivo*, in adipose tissue of high fat fed obese mice [86]. Despite the lack of detectable difference in adipose tissue macrophage phenotype, an increased angiogenic response to tissue ischemia is also seen in adipose tissue of obese *Hsd11b1*<sup>-/-</sup> mice and underlies their resistance to some of the adverse metabolic consequences of obesity [87], suggesting the pro-angiogenic phenotype may be at least partly independent of macrophages.

How is the improved recovery of *Hsd11b1*<sup>-/-</sup> mice from inflammation following myocardial infarction reconciled with our original hypothesis? As predicted by the hypothesis, deficiency in 11 $\beta$ -HSD1 causes greater release of pro-inflammatory cytokines from LPS-treated macrophages [1, 81], suggesting an exaggerated M1 macrophage phenotype. However, the earlier switch to an M2 phenotype was unexpected. Whether this reflects a switch to M2 phenotype *in situ* or recruitment of a distinct subset of monocytes is currently unknown. It is possible that this is a consequence of prolonged activation of the HPA axis in *Hsd11b1*<sup>-/-</sup> mice. However, these mice show little perturbation of plasma corticosterone levels, even following



stress, on this genetic background [88], so the earlier switch is unlikely to be mediated by plasma glucocorticoids. Moreover, as discussed above, intracellular amplification of glucocorticoid signalling by 11 $\beta$ -HSD1 is predicted to accelerate repair and resolution processes, not attenuate them. Several key factors implicated in macrophage polarisation [89] are differentially expressed in *Hsd11b1*<sup>-/-</sup> mice. The Src homology 2-containing inositol-5'-phosphatase (SHIP)-1 negatively regulates the PI3K pathway. It represses the generation of M2 macrophages [74] yet restrains LPS-induced (M1) activation of bone marrow-derived (naïve) macrophages [90]. Moreover, elevated SHIP1 expression induces endotoxin tolerance [90] with reduced pro-inflammatory cytokine production with subsequent endotoxin challenge [90]. The increased LPS-responsiveness of thioglycollate elicited peritoneal (TEP) macrophages from *Hsd11b1*<sup>-/-</sup> mice was attributed to elevated SHIP1 levels as a consequence of higher levels of TGF $\beta$ 1 [81] though SHIP1 levels appear to decrease more rapidly following LPS in *Hsd11b1*<sup>-/-</sup> macrophages than in wild-type. In splenic macrophages, basal SHIP1 levels are normal in *Hsd11b1*<sup>-/-</sup> mice, but unlike wild-type splenic macrophages, those from *Hsd11b1*<sup>-/-</sup> mice fail to down-regulate SHIP1 following LPS [81]. Whether this induces endotoxin tolerance [90] to a greater extent in *Hsd11b1*<sup>-/-</sup> macrophages is something that requires testing. Thus, SHIP1 appears abnormally regulated in *Hsd11b1*<sup>-/-</sup> macrophages, though why is currently unclear. Nevertheless, these somewhat confusing data illustrate that M1/M2 macrophage polarisation in *Hsd11b1*<sup>-/-</sup> mice may be highly dependent upon the macrophage population and context.

Hypoxia-inducible factor (HIF1)- $\alpha$ , which promotes M1 polarisation, is decreased in adipose tissue of *Hsd11b1*<sup>-/-</sup> mice, whereas levels of PPAR $\alpha$  (which promotes the M2 phenotype) are increased [87, 91]. Whether these factors are differentially expressed in macrophages of *Hsd11b1*<sup>-/-</sup> mice will be important to determine.

The outcome of acute inflammation is not invariably improved in *Hsd11b1*<sup>-/-</sup> mice. At the stage when arthritis has largely resolved in wild-type mice following K/BxN serum transfer, joints of *Hsd11b1*<sup>-/-</sup> mice show greater periarticular fibrosis, more extensive exostoses and ganglion cyst formation. Following carageenan-induced pleurisy, *Hsd11b1*<sup>-/-</sup> mice show persistence of inflammation at a stage when it is resolving in wild-type controls, as well as lymphoid aggregates within the lung and formation of fibrous adhesions between lung lobes, the latter not present in control mice [83]. Whether these disadvantageous features result from greater inflammation in *Hsd11b1*<sup>-/-</sup> mice, an earlier switch to a pro-repair (pro-fibrotic) M2 phenotype, a greater response of the non-immune tissue or a combination of all of these will be an interesting question for the future. Moreover, the consequences of more extended inflammation will be interesting to determine. The preliminary findings in arthritis and carageenan induced pleurisy suggest that 11 $\beta$ -HSD1-deficiency or inhibition may aggravate diseases associated with a dysregulated angiogenic and pro-fibrotic phenotype, including rheumatoid arthritis.

### *Chronic inflammation*

Chronic inflammation results from a failure to resolve acute inflammation. Atherosclerosis, diabetes, metabolic syndrome and Alzheimer's disease are all now recognised as chronic inflammatory diseases. Even simple obesity is frequently associated with low level chronic inflammation within the adipose tissue. The elevation in systemic pro-inflammatory cytokines during chronic inflammation might be expected to activate the HPA axis. However, plasma cortisol is normal in both the "classic" inflammatory diseases (rheumatoid arthritis, inflammatory bowel disease etc) and in the "metabolic" inflammatory diseases (atherosclerosis, metabolic syndrome, diabetes), at least until these become complicated by additional pathologies. HPA axis activity may be elevated in metabolic inflammation, with increased clearance of

glucocorticoids maintaining normal plasma cortisol levels [92] but possibly increasing plasma cortisone levels (and thus 11 $\beta$ -HSD1 substrate), though this has only been indirectly measured. In rheumatoid arthritis and other inflammatory diseases however, the HPA axis appears relatively suppressed, especially given the level of systemic inflammation expected to activate the axis [4, 93]. Edwards has recently hypothesised that this apparent deficiency in HPA activation is a result of the systemic increase in TNF- $\alpha$  in chronic inflammation inducing a widespread increase in 11 $\beta$ -HSD1 expression, including in the hypothalamus, thus amplifying negative feedback by glucocorticoids on the HPA axis [93]. Whether this is indeed the case requires experimental testing, but consistent with this hypothesis, whole body conversion of cortisone to cortisol (relative to cortisol to cortisone) is increased in patients with inflammatory disease [94] suggesting altered balance of 11 $\beta$ -HSD activities in favour of 11 $\beta$ -reductase (11 $\beta$ -HSD1).

#### *Metabolic Syndrome, type 2 diabetes and atherosclerosis*

11 $\beta$ -HSD1 deficiency or inhibition is metabolically beneficial in rodent models of diet-induced obesity or diabetes. It improves hepatic and adipose insulin sensitivity, attenuates hepatic gluconeogenesis, skews to a “cardioprotective” plasma lipid profile, shifts hepatic lipid metabolism from lipogenesis to fatty acid oxidation and causes a preferential gain of peripheral adipose tissue at the expense of visceral [86, 91, 95-101] (and see Figure 2). Similarly, in patients with type 2 diabetes, 11 $\beta$ -HSD1 inhibition lowers plasma glucose and lipids, consistent with rodent studies. It also modestly reduces blood pressure in human hypertension [102-104]. Intriguingly, an 11 $\beta$ -HSD1 inhibitor more effectively improved glucose homeostasis in obese mice when administered close to the time of the diurnal peak of plasma glucocorticoid levels [105]. Given that 11 $\beta$ -HSD1 mRNA probably does not vary with the circadian rhythm [105, 106] (though one study suggests it may in rats [107]), this is much more likely to reflect high 11 $\beta$ -HSD1 substrate levels at peak HPA axis activity [108]. Indeed, 11 $\beta$ -HSD1 may contribute to normal circadian control of the HPA axis, at least in some genetic backgrounds [88, 108]. 11 $\beta$ -HSD1 is expressed in the paraventricular nucleus of the human hypothalamus, suggesting a conserved role in HPA axis regulation [109].

Recent data suggest that the liver is not the sole or even predominant target of the metabolically beneficial effects of 11 $\beta$ -HSD1-deficiency or inhibition; conditional deletion of 11 $\beta$ -HSD1 in hepatocytes of mice produces only minimal improvements in glucose homeostasis in diet-induced obesity [110]. Instead, increased glucocorticoid activity in adipose tissue is implicated. In obese humans, numerous studies have reported elevated 11 $\beta$ -HSD1 expression in subcutaneous adipose tissue (reviewed [92]) and in human omental fat, 11 $\beta$ -HSD1 expression correlates with adipocyte hypertrophy [111, 112], itself associated with a more pro-inflammatory state [113, 114]. In mice, a two to three-fold elevation of 11 $\beta$ -HSD1 selectively in adipose tissue phenocopies the metabolic syndrome, with central obesity, insulin resistance, dyslipidaemia and hypertension [115, 116] whereas similar transgenic expression of 11 $\beta$ -HSD2 in adipocytes (it is not normally expressed in adipocytes), presumably lowering intra-adipose glucocorticoid action, causes insulin sensitisation in high fat fed mice [117].

11 $\beta$ -HSD1-deficiency protects against pro-inflammatory changes in adipose tissue in obesity. Inflammatory cell (macrophages, lymphocytes) infiltration of mesenteric adipose tissue is lower in high fat-fed 11 $\beta$ -HSD1-deficient mice than in controls, probably due to reduced adipocyte secretion of the pro-inflammatory chemokine, monocyte chemoattractant-1 (MCP-1) [86]. This is associated with higher levels of AMP-activated protein kinase activation in this depot [86], likely to contribute to the maintained lipid oxidation with obesity [118] in 11 $\beta$ -HSD1-deficiency. Whether these

changes are a cause or a consequence of the increase in angiogenesis and reduction in hypoxia and fibrosis recently described in the adipose tissue of these mice [87] is an interesting question. Adipose tissue hypoxia is associated with a local pro-inflammatory environment and leads to fibrosis though not necessarily angiogenesis [119-121], suggesting that it is the greater angiogenic response in *Hsd11b1*<sup>-/-</sup> mice that is protective against adipose tissue hypoxia and fibrosis. PPAR $\alpha$  mRNA levels are higher and the pro-angiogenic response to PPAR $\alpha$  activation is much greater in *Hsd11b1*<sup>-/-</sup> adipocytes than in controls, placing the adipocyte at the heart of the response. Whether there are also beneficial roles for macrophage and/or vascular 11 $\beta$ -HSD1 is important to determine.

As well as improving metabolic risk factors, deficiency in or inhibition of 11 $\beta$ -HSD1 also reduces atherosclerosis and systemic inflammation and lowers macrophage and T cell infiltration of atherosclerotic lesions in *Apoe*<sup>-/-</sup> mice [122-124]. This is the converse of what happens with 11 $\beta$ -HSD2-deficiency, which is pro-inflammatory in the endothelium and accelerates atherosclerosis in *Apoe*<sup>-/-</sup> mice, an effect at least partly mediated through activation of the MR as it is blocked by eplerenone, an MR antagonist [125]. The atheroprotective effects of 11 $\beta$ -HSD1-deficiency are likely to be mediated through both systemic (reduced circulating monocyte chemotactic protein (MCP)-1 and number of pro-inflammatory Ly6C<sup>hi</sup> monocytes) and local (reduced aortic vascular cell adhesion molecule (VCAM)-1 expression) mechanisms [124]. It is interesting to speculate that reduced visceral adipose tissue inflammation may contribute to the reduction in systemic inflammation – as in diet-induced obesity, mesenteric adipose tissue MCP-1 mRNA levels are reduced in western diet-fed 11 $\beta$ -HSD1-deficient *Apoe*<sup>-/-</sup> mice [124].

#### *“Classic” inflammatory diseases - rheumatoid arthritis*

If 11 $\beta$ -HSD1-deficiency is beneficial in chronic “cardiometabolic inflammation”, what of the classical inflammatory diseases, in which a glucocorticoid-insufficient state is suggested and glucocorticoid therapy remains highly effective? Inevitably, studies in animals are predominantly short term, modelling the disease, whereas the disease in patients frequently reflects years of accumulated damage and inflammation. These situations may be quite different. Nevertheless, accumulating evidence in both patients and animal models is consistent with dysregulated 11 $\beta$ -HSD1 in the inflamed joint in rheumatoid arthritis as well as increased colonic expression of 11 $\beta$ -HSD1 at sites of inflammation in inflammatory bowel disease (reviewed [35]). So far, studies in inflammatory bowel disease have gone little beyond observation, though they do suggest that at least some of the increase in 11 $\beta$ -HSD1 expression occurs in activated lymphocytes that migrate from the inflamed colon to the draining lymph nodes [54]. Studies in human patients with rheumatoid arthritis suggest differential regulation of 11 $\beta$ -HSDs in immune and mesenchymal cells. Comparison of cortisone and cortisol levels in synovial fluid and serum suggest the balance favours intra-articular generation of cortisol in the rheumatic joint [126] although it seems that even so, the overall capacity to convert cortisone to cortisol may be reduced in the inflamed arthritic synovium compared to non-inflamed. However, within inflamed rheumatic joints, synovial inflammation still correlates with conversion of cortisone to cortisol [127]. This complex relationship probably reflects the balance between high expression of 11 $\beta$ -HSD1 in synovial fibroblasts from arthritic patients (almost certainly as a result of the pro-inflammatory cytokine environment) and expression of 11 $\beta$ -HSD2 in synovial macrophages from patients with rheumatoid arthritis [126-128]. This latter finding accords with other studies identifying 11 $\beta$ -HSD2 as a peripheral blood mononuclear cell marker of early rheumatoid arthritis and highly expressed in the arthritic joint [129, 130]. 11 $\beta$ -HSD2-positive macrophages have also been described in the lungs of patients who died of acute respiratory distress syndrome [131]. Similar cells

(macrophages, lymphocytes) from healthy humans do not express 11 $\beta$ -HSD2 [37, 130], nor has 11 $\beta$ -HSD2 been found in mouse leukocytes [1]. 11 $\beta$ -HSD2 expression in leukocytes may reflect a species difference between mouse and human, or could, in humans, reflect an adaptive response to chronic inflammation. The biological reason for this apparently pro-inflammatory change is unknown but it is likely to cause resistance to endogenous glucocorticoids, which might be overcome by pharmacological levels of synthetic glucocorticoids like prednisolone or bypassed with non-metabolised synthetic glucocorticoids like dexamethasone.

What might 11 $\beta$ -HSD1 inhibition do in chronic inflammatory disease? If the Edwards hypothesis [93] is correct, then systemic inhibition of 11 $\beta$ -HSD1, particularly if administered during the night (in humans), should correct the HPA axis abnormality and boost the plasma cortisol levels. This might be enough to dampen down some of the inflammation, though 11 $\beta$ -HSD1 inhibition would also deprive inflamed tissues of the 11 $\beta$ -HSD1-mediated increase in intracellular glucocorticoid levels. Moreover, given that cortisol also activates MR (in the absence of 11 $\beta$ -HSD2), this could further exacerbate inflammation which could be particularly damaging within the vasculature (see below). In chronic inflammatory disease, continuing tissue injury is frequently associated with fibrosis and angiogenesis. Both may be exacerbated by 11 $\beta$ -HSD1 inhibition. As mentioned above, 11 $\beta$ -HSD1-deficient mice show an increased angiogenic response to adipose tissue hypoxia, to ischaemia following myocardial infarction, in wound healing and in sub-cutaneously implanted sponges [85, 87]. They also show a pro-fibrotic response to pleural inflammation and following inflammatory arthritis [83]. Whilst it is currently unclear whether the increased fibrosis in 11 $\beta$ -HSD1-deficient mice will resolve completely during recovery from inflammation, it is likely that if the injurious stimulus persists, fibrosis will be more severe with 11 $\beta$ -HSD1-deficiency or inhibition. In continuing liver injury, a population of macrophages with "M2"-like properties drives the fibrotic response, probably mediated at least in part through TGF $\beta$  1 [40]. Higher macrophage expression of TGF $\beta$  1 with 11 $\beta$ -HSD1-deficiency [81] may be an important contributor to the pro-fibrotic phenotype of these mice.

#### *Glucocorticoid receptor or mineralocorticoid receptor activation?*

Activation of MR, most notably in the heart and vasculature, has pro-inflammatory and pro-fibrotic consequences [132, 133]. Unlike synthetic glucocorticoids, most of which show selectivity for GR over MR, endogenous glucocorticoids bind with higher affinity to MR than to GR. Thus, MR is usually considered near saturated at circulating glucocorticoid levels, even at the diurnal nadir [134]. Aldosterone activates MR irrespective of which cells it is expressed in, but cortisol activation of MR is normally prevented if 11 $\beta$ -HSD2 is co-expressed with MR. However, under conditions of oxidative stress, endogenous glucocorticoids can activate MR, at least in the cardiovascular system [135]. A crucial question therefore, central to the function of 11 $\beta$ -HSD1, is which receptor binds the ligand it generates, GR or MR? This may differ according to tissues. MR is absent from liver, so in this tissue, 11 $\beta$ -HSD1 provides ligand to GR. However, MR is expressed in some classical glucocorticoid targets, including adipocytes and macrophages, normally in the absence of 11 $\beta$ -HSD2, where it presumably functions as a glucocorticoid receptor. A pro-inflammatory role for glucocorticoid-activated MR is suggested; eplerenone treatment of *ob/ob* mice prevented the obesity-associated increases in MCP-1, TNF- $\alpha$  and other inflammatory markers in adipose tissue [136]. Whether the relevant cell is the adipocyte, however, is unclear. Whereas MR activation (presumably by glucocorticoids) in macrophages appears pro-inflammatory, macrophage-specific deletion of MR appears anti-inflammatory - it causes M2 polarisation of macrophages [43] and reduces cerebral

infarct area following ischaemia in mice, concomitant with reduced expression of M1 macrophage markers (TNF- $\alpha$ , IL-1, MCP-1 etc) but maintained M2 markers (Ym1, Arg1) [137]. Thus, the consequences of 11 $\beta$ -HSD1-mediated glucocorticoid generation could differ greatly, depending on cellular oxidation/stress state and the relative levels of GR *versus* MR.

## SUMMARY AND CONCLUSIONS

Consistent with the adverse metabolic effects of glucocorticoid excess, 11 $\beta$ -HSD1 deficiency or inhibition is clearly beneficial in cardiometabolic disease. The extent to which this is dependent on inhibition/deficiency within inflammatory cells will be interesting to discover. Also, whether 11 $\beta$ -HSD1 deficiency/inhibition is beneficial in other types of inflammation remains to be seen. Current evidence suggests that the acute response to injury is more severe. The subsequent recovery phase may depend on whether the injurious stimulus persists as in patients with rheumatoid arthritis (in which case 11 $\beta$ -HSD1 deficiency/inhibition may worsen the disease), or whether recovery and tissue remodelling occur, as for example follows myocardial infarction (when 11 $\beta$ -HSD1 deficiency/inhibition may aid recovery). The application of Cre/Lox technology to generate tissue- and cell-specific “knock-out” of 11 $\beta$ -HSD1 will be invaluable in dissecting the contributions of immune cells, particularly macrophages and neutrophils, to the pro-angiogenic and pro-fibrotic phenotype. In the future, such studies could lead to better targeting of glucocorticoid therapy, perhaps even targeting macrophages separately from host tissues at specific temporal stages of disease. As already suggested [1], targeted delivery of inactive glucocorticoid precursors to macrophages might provide an effective future therapy for chronic inflammatory disease.

## ACKNOWLEDGEMENTS

Work in the authors’ laboratory is funded by grants from the Wellcome Trust and the Medical Research Council. We thank colleagues in the University/BHF Centre for Cardiovascular Science and the MRC Centre for Inflammation Research for many stimulating discussions.

## REFERENCES

- [1] J.S. Gilmour, A.E. Coutinho, J.F. Cailhier, T.Y. Man, M. Clay, G. Thomas, H.J. Harris, J.J. Mullins, J.R. Seckl, J.S. Savill, K.E. Chapman, Local amplification of glucocorticoids by 11 $\beta$ -hydroxysteroid dehydrogenase type 1 promotes macrophage phagocytosis of apoptotic leukocytes, *J Immunol* 176 (2006) 7605-7611.
- [2] J.A. Brewer, B. Khor, S.K. Vogt, L.M. Muglia, H. Fujiwara, K.E. Haegele, B.P. Sleckman, L.J. Muglia, T-cell glucocorticoid receptor is required to suppress COX-2-mediated lethal immune activation, *Nat Med* 9 (2003) 1318-1322.
- [3] S. Bhattacharyya, D.E. Brown, J.A. Brewer, S.K. Vogt, L.J. Muglia, Macrophage glucocorticoid receptors regulate Toll-like receptor-4-mediated inflammatory responses by selective inhibition of p38 MAP kinase, *Blood* 109 (2007) 4313-4319.
- [4] G.P. Chrousos, The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation, *N Engl J Med* 332 (1995) 1351-1362.
- [5] A.V. Turnbull, C.L. Rivier, Regulation of the hypothalamic-pituitary-adrenal axis by cytokines: actions and mechanisms of action, *Physiol Rev* 79 (1999) 1-71.
- [6] M.S. Harbuz, A.J. Chover-Gonzalez, D.S. Jessop, Hypothalamo-pituitary-adrenal axis and chronic immune activation, *Ann N Y Acad Sci* 992 (2003) 99-106.

- 578 [7] M. Cutolo, A. Sulli, C. Pizzorni, M.E. Secchi, S. Soldano, B. Seriolo, R.H. Straub,  
579 K. Otsa, G.J. Maestroni, Circadian rhythms: glucocorticoids and arthritis, *Ann N Y*  
580 *Acad Sci* 1069 (2006) 289-299.
- 581 [8] J. Savill, I. Dransfield, C. Gregory, C. Haslett, A blast from the past: clearance of  
582 apoptotic cells regulates immune responses, *Nat Rev Immunol* 2 (2002) 965-975.
- 583 [9] D.W. Gilroy, The endogenous control of acute inflammation - from onset to  
584 resolution, *Drug Discovery Today: Therapeutic Strategies* 1 (2004) 313.
- 585 [10] C.N. Serhan, J. Savill, Resolution of inflammation: the beginning programs the  
586 end, *Nat Immunol* 6 (2005) 1191-1197.
- 587 [11] P.J. Barnes, Anti-inflammatory actions of glucocorticoids: molecular mechanisms,  
588 *Clin Sci (Lond)* 94 (1998) 557-572.
- 589 [12] K.A. Smoak, J.A. Cidlowski, Mechanisms of glucocorticoid receptor signaling  
590 during inflammation, *Mech Ageing Dev* 125 (2004) 697-706.
- 591 [13] M. Perretti, A. Ahluwalia, The microcirculation and inflammation: site of action  
592 for glucocorticoids, *Microcirculation* 7 (2000) 147-161.
- 593 [14] B.S. McEwen, C.A. Biron, K.W. Brunson, K. Bulloch, W.H. Chambers, F.S.  
594 Dhabhar, R.H. Goldfarb, R.P. Kitson, A.H. Miller, R.L. Spencer, J.M. Weiss, The  
595 role of adrenocorticoids as modulators of immune function in health and disease:  
596 neural, endocrine and immune interactions, *Brain Res Rev* 23 (1997) 79-133.
- 597 [15] S.L. Planey, G. Litwack, Glucocorticoid-induced apoptosis in lymphocytes,  
598 *Biochem Biophys Res Commun* 279 (2000) 307-312.
- 599 [16] J.D. Ashwell, F.W. Lu, M.S. Vacchio, Glucocorticoids in T cell development and  
600 function, *Ann Rev Immunol* 18 (2000) 309-345.
- 601 [17] M.J. Herold, K.G. McPherson, H.M. Reichardt, Glucocorticoids in T cell apoptosis  
602 and function, *Cell Mol Life Sci* 63 (2006) 60-72.
- 603 [18] A. McColl, S. Michlewska, I. Dransfield, A.G. Rossi, Effects of glucocorticoids on  
604 apoptosis and clearance of apoptotic cells, *ScientificWorldJournal* 7 (2007) 1165-  
605 1181.
- 606 [19] A.E. Coutinho, K.E. Chapman, The anti-inflammatory and immunosuppressive  
607 effects of glucocorticoids, recent developments and mechanistic insights, *Mol Cell*  
608 *Endocrinol* 335 (2011) 2-13.
- 609 [20] K.M. Giles, K. Ross, A.G. Rossi, N.A. Hotchin, C. Haslett, I. Dransfield,  
610 Glucocorticoid augmentation of macrophage capacity for phagocytosis of apoptotic  
611 cells is associated with reduced p130Cas expression, loss of paxillin/pyk2  
612 phosphorylation, and high levels of active Rac, *J Immunol* 167 (2001) 976-986.
- 613 [21] J. Ehrchen, L. Steinmuller, K. Barczyk, K. Tenbrock, W. Nacken, M. Eisenacher,  
614 U. Nordhues, C. Sorg, C. Sunderkotter, J. Roth, Glucocorticoids induce  
615 differentiation of a specifically activated, anti-inflammatory subtype of human  
616 monocytes, *Blood* 109 (2007) 1265-1274.
- 617 [22] F. Vallelleian, C.A. Schaer, T. Kaempfer, P. Gehrig, E. Duerst, G. Schoedon, D.J.  
618 Schaer, Glucocorticoid treatment skews human monocyte differentiation into a  
619 hemoglobin-clearance phenotype with enhanced heme-iron recycling and  
620 antioxidant capacity, *Blood* 116 (2011) 5347-5356.
- 621 [23] A. Tsianakas, G. Varga, K. Barczyk, G. Bode, N. Nippe, N. Kran, J. Roth, T.A.  
622 Luger, J. Ehrchen, C. Sunderkoetter, Induction of an anti-inflammatory human  
623 monocyte subtype is a unique property of glucocorticoids, but can be modified by  
624 IL-6 and IL-10, *Immunobiology* 217 (2012) 329-335.
- 625 [24] G. Varga, J. Ehrchen, A. Tsianakas, K. Tenbrock, A. Rattenholl, S. Seeliger, M.  
626 Mack, J. Roth, C. Sunderkoetter, Glucocorticoids induce an activated, anti-

- inflammatory monocyte subset in mice that resembles myeloid-derived suppressor cells, *J Leukoc Biol* 84 (2008) 644-650.
- [25] S. Schif-Zuck, N. Gross, S. Assi, R. Rostoker, C.N. Serhan, A. Ariel, Saturated-efferocytosis generates pro-resolving CD11b low macrophages: Modulation by resolvins and glucocorticoids, *Eur J Immunol* 41 (2011) 366-379.
- [26] M.P. Yeager, P.M. Guyre, A.U. Munck, Glucocorticoid regulation of the inflammatory response to injury, *Acta Anaesthesiol Scand* 48 (2004) 799-813.
- [27] H.Y. Lim, N. Muller, M.J. Herold, J. van den Brandt, H.M. Reichardt, Glucocorticoids exert opposing effects on macrophage function dependent on their concentration, *Immunology* 122 (2007) 47-53.
- [28] M. Lombes, S. Kenouch, A. Souque, N. Farman, M.E. Rafestin-Oblin, The mineralocorticoid receptor discriminates aldosterone from glucocorticoids independently of the 11 $\beta$ -hydroxysteroid dehydrogenase, *Endocrinology* 135 (1994) 834-840.
- [29] A.H. Miller, R.L. Spencer, M. Stein, B.S. McEwen, Adrenal steroid receptor binding in spleen and thymus after stress or dexamethasone, *Am J Physiol* 259 (1990) E405-E412.
- [30] G.D. Barish, M. Downes, W.A. Alaynick, R.T. Yu, C.B. Ocampo, A.L. Bookout, D.J. Mangelsdorf, R.M. Evans, A Nuclear Receptor Atlas: macrophage activation, *Mol Endocrinol* 19 (2005) 2466-2477.
- [31] S. Diederich, E. Eigendorff, P. Burkhardt, M. Quinkler, C. Bumke-Vogt, M. Rochel, D. Seidelmann, P. Esperling, W. Oelkers, V. Bahr, 11 $\beta$ -Hydroxysteroid dehydrogenase types 1 and 2: an important pharmacokinetic determinant for the activity of synthetic mineralo- and glucocorticoids, *J Clin Endocrinol Metab* 87 (2002) 5695-5701.
- [32] J.R. Seckl, 11 $\beta$ -hydroxysteroid dehydrogenases: changing glucocorticoid action., *Curr Opin Pharmacol* 4 (2004) 597-602.
- [33] C.R.W. Edwards, P.M. Stewart, D. Burt, L. Brett, M.A. McIntyre, W.S. Sutanto, E.R. de Kloet, C. Monder, Localisation of 11 $\beta$ -hydroxysteroid dehydrogenase-tissue specific protector of the mineralocorticoid receptor, *Lancet* ii (1988) 986-989.
- [34] J.W. Funder, P.T. Pearce, R. Smith, A.I. Smith, Mineralocorticoid action: target tissue specificity is enzyme, not receptor, mediated, *Science* 242 (1988) 583-585.
- [35] K.E. Chapman, A.E. Coutinho, M. Gray, J.S. Gilmour, J.S. Savill, J.R. Seckl, The role and regulation of 11 $\beta$ -hydroxysteroid dehydrogenase type 1 in the inflammatory response., *Mol Cell Endocrinol* 301 (2009) 123-131.
- [36] A.E. Coutinho, T. Kipari, Z. Zhang, C. Esteves, J.S. Gilmour, J.-F. Cailhier, J. Hughes, J.R. Seckl, J.S. Savill, K.E. Chapman, Dynamic Regulation of 11 $\beta$ -Hydroxysteroid Dehydrogenase Type 1 in Neutrophils during an Inflammatory Response, *Endocrine Reviews* 32 (2011) P2-575.
- [37] R. Thieringer, C.B. Le Grand, L. Carbin, T.Q. Cai, B. Wong, S.D. Wright, A. Hermanowski-Vosatka, 11 $\beta$ -Hydroxysteroid dehydrogenase type 1 is induced in human monocytes upon differentiation to macrophages, *J Immunol* 167 (2001) 30-35.
- [38] F.O. Martinez, A. Sica, A. Mantovani, M. Locati, Macrophage activation and polarization, *Front Biosci* 13 (2008) 453-461.
- [39] F.O. Martinez, L. Helming, S. Gordon, Alternative activation of macrophages: an immunologic functional perspective, *Annu Rev Immunol* 27 (2009) 451-483.

- 675 [40] J.S. Duffield, S.J. Forbes, C.M. Constandinou, S. Clay, M. Partolina, S. Vuthoori,  
676 S. Wu, R. Lang, J.P. Iredale, Selective depletion of macrophages reveals distinct,  
677 opposing roles during liver injury and repair, *J Clin Invest* 115 (2005) 56-65.
- 678 [41] Y.Q. Liu, J.M. Cousin, J. Hughes, J. VanDamme, J.R. Seckl, C. Haslett, I.  
679 Dransfield, J. Savill, A.G. Rossi, Glucocorticoids promote nonphlogistic  
680 phagocytosis of apoptotic leukocytes, *J Immunol* 162 (1999) 3639-3646.
- 681 [42] A. Kleiman, S. Hubner, J.M. Rodriguez Parkitna, A. Neumann, S. Hofer, M.A.  
682 Weigand, M. Bauer, W. Schmid, G. Schutz, C. Libert, H.M. Reichardt, J.P.  
683 Tuckermann, Glucocorticoid receptor dimerization is required for survival in septic  
684 shock via suppression of interleukin-1 in macrophages, *FASEB J* 26 (2012) 722-  
685 729.
- 686 [43] M.G. Usher, S.Z. Duan, C.Y. Ivaschenko, R.A. Frieler, S. Berger, G. Schutz, C.N.  
687 Lumeng, R.M. Mortensen, Myeloid mineralocorticoid receptor controls  
688 macrophage polarization and cardiovascular hypertrophy and remodeling in mice,  
689 *The Journal of Clinical Investigation* 120 (2010) 3350-3364.
- 690 [44] J.S. Gilmour, Glucocorticoids, 11 $\beta$ -hydroxysteroid dehydrogenases and  
691 macrophage function., PhD thesis, University of Edinburgh, 2003.
- 692 [45] F.O. Martinez, S. Gordon, M. Locati, A. Mantovani, Transcriptional profiling of  
693 the human monocyte-to-macrophage differentiation and polarization: new  
694 molecules and patterns of gene expression, *J Immunol* 177 (2006) 7303-7311.
- 695 [46] V. Joganathan, A. Al-Hakami, S. Rauz, P.M. Stewart, G.R. Wallace, I.J. Bujalska,  
696 Local cortisol generation by human macrophage subsets by 11 $\beta$ -hydroxysteroid  
697 dehydrogenase type 1 enzyme and its role in ocular immune privilege, *Endocrine*  
698 *Abstracts* 15 (2008) OC30.
- 699 [47] G. Chinetti-Gbaguidi, M.A. Bouhrel, C. Copin, C. Duhem, B. Derudas, B. Neve, B.  
700 Noel, J. Eeckhoutte, P. Lefebvre, J.R. Seckl, B. Staels, Peroxisome Proliferator  
701 Activated Receptor- $\alpha$  Activation Induces 11 $\beta$ -Hydroxysteroid Dehydrogenase  
702 Type 1 Activity in Human Alternative Macrophages, *Arterioscler Thromb Vasc*  
703 *Biol* 32 (2012) 677-685.
- 704 [48] J.C. Rodriguez-Prados, P.G. Traves, J. Cuenca, D. Rico, J. Aragonés, P. Martín-  
705 Sanz, M. Cascante, L. Bosca, Substrate fate in activated macrophages: a  
706 comparison between innate, classic, and alternative activation, *J Immunol* 185  
707 (2010) 605-614.
- 708 [49] A. Haschemi, P. Kosma, L. Gille, C.R. Evans, C.F. Burant, P. Starkl, B. Knapp, R.  
709 Haas, J.A. Schmid, C. Jandl, S. Amir, G. Lubec, J. Park, H. Esterbauer, M. Bilban,  
710 L. Brizuela, J.A. Pospisilik, L.E. Otterbein, O. Wagner, The Sedoheptulose Kinase  
711 CARKL Directs Macrophage Polarization through Control of Glucose Metabolism,  
712 *Cell Metab* 15 (2012) 813-826.
- 713 [50] V.A. Fadok, D.L. Bratton, A. Konowal, P.W. Freed, J.Y. Westcott, P.M. Henson,  
714 Macrophages that have ingested apoptotic cells in vitro inhibit proinflammatory  
715 cytokine production through autocrine/paracrine mechanisms involving TGF- $\beta$ ,  
716 PGE<sub>2</sub>, and PAF, *J Clin Invest* 101 (1998) 890-898.
- 717 [51] K.E. Chapman, A. Coutinho, M. Gray, J.S. Gilmour, J.S. Savill, J.R. Seckl, Local  
718 amplification of glucocorticoids by 11 $\beta$ -hydroxysteroid dehydrogenase type 1 and  
719 its role in the inflammatory response, *Ann N Y Acad Sci* 1088 (2006) 265-273.
- 720 [52] J. Bryndova, S. Zbankova, M. Kment, J. Pacha, Colitis up-regulates local  
721 glucocorticoid activation and down-regulates inactivation in colonic tissue, *Scand J*  
722 *Gastroenterol* 39 (2004) 549-553.



- [53] P. Ergang, P. Leden, K. Vagnerova, P. Klusonova, I. Miksik, J. Jurcovicova, M. Kment, J. Pacha, Local metabolism of glucocorticoids and its role in rat adjuvant arthritis, *Mol Cell Endocrinol* 323 (2010) 155-160.
- [54] P. Ergang, K. Vytackova, J. Svec, J. Bryndova, I. Miksik, J. Pacha, Upregulation of 11 $\beta$ -hydroxysteroid dehydrogenase 1 in lymphoid organs during inflammation in the rat, *J Steroid Biochem Mol Biol* 126 (2011) 19-25.
- [55] A.R. Dover, P.W. Hadoke, L.J. Macdonald, E. Miller, D.E. Newby, B.R. Walker, Intravascular glucocorticoid metabolism during inflammation and injury in mice, *Endocrinology* 148 (2007) 166-172.
- [56] Z. Yang, X. Zhu, C. Guo, K. Sun, Stimulation of 11 $\beta$ -HSD1 expression by IL-1 $\beta$  via a C/EBP binding site in human fetal lung fibroblasts, *Endocrine* 36 (2009) 404-411.
- [57] I.D. Ignatova, R.M. Kostadinova, C.E. Goldring, A.R. Nawrocki, F.J. Frey, B.M. Frey, Tumor necrosis factor- $\alpha$  upregulates 11 $\beta$ -hydroxysteroid dehydrogenase type 1 expression by CCAAT/enhancer binding protein-1 in HepG2 cells, *Am J Physiol Endocrinol Metab* 296 (2009) E367-377.
- [58] M.M. Ahasan, R. Hardy, C. Jones, K. Kaur, D. Nanus, M. Juarez, S.A. Morgan, Z. Hassan-Smith, C. Benezech, J.H. Caamano, M. Hewison, G. Lavery, E.H. Rabbitt, A.R. Clark, A. Filer, C.D. Buckley, K. Raza, P.M. Stewart, M.S. Cooper, Inflammatory regulation of glucocorticoid metabolism in mesenchymal stromal cells, *Arthritis Rheum* 64 (2012) 2404-2413.
- [59] R.M. Kostadinova, A.R. Nawrocki, F.J. Frey, B.M. Frey, Tumor necrosis factor  $\alpha$  and phorbol 12-myristate-13-acetate down-regulate human 11 $\beta$ -hydroxysteroid dehydrogenase type 2 through p50/p50 NF-kappaB homodimers and Egr-1, *FASEB J* 19 (2005) 650-652.
- [60] K. Sun, L. Myatt, Enhancement of glucocorticoid-induced 11 $\beta$ -hydroxysteroid dehydrogenase type 1 expression by proinflammatory cytokines in cultured human amnion fibroblasts, *Endocrinology* 144 (2003) 5568-5577.
- [61] M.T. Rae, D. Niven, H.O. Critchley, C.R. Harlow, S.G. Hillier, Antiinflammatory steroid action in human ovarian surface epithelial cells, *J Clin Endocrinol Metab* 89 (2004) 4538-4544.
- [62] W. Li, L. Gao, Y. Wang, T. Duan, L. Myatt, K. Sun, Enhancement of cortisol-induced 11 $\beta$ -hydroxysteroid dehydrogenase type 1 expression by interleukin 1 $\beta$  in cultured human chorionic trophoblast cells, *Endocrinology* 147 (2006) 2490-2495.
- [63] K. Kaur, R. Hardy, M.M. Ahasan, M. Eijken, J.P. van Leeuwen, A. Filer, A.M. Thomas, K. Raza, C.D. Buckley, P.M. Stewart, E.H. Rabbitt, M. Hewison, M.S. Cooper, Synergistic induction of local glucocorticoid generation by inflammatory cytokines and glucocorticoids: implications for inflammation associated bone loss, *Ann Rheum Dis* 69: (2010) 1185-1190.
- [64] L.J.S. Williams, V. Lyons, I. MacLeod, V. Rajan, G.J. Darlington, V. Poli, J.R. Seckl, K.E. Chapman, C/EBP regulates hepatic transcription of 11 $\beta$ -hydroxysteroid dehydrogenase type 1; a novel mechanism for cross talk between the C/EBP and glucocorticoid signalling pathways, *J Biol Chem* 275 (2000) 30232-30239.
- [65] J. Gout, J. Tirard, C. Thevenon, J.P. Riou, M. Begeot, D. Naville, CCAAT/enhancer-binding proteins (C/EBPs) regulate the basal and cAMP-induced transcription of the human 11 $\beta$ -hydroxysteroid dehydrogenase encoding gene in adipose cells, *Biochimie* 88 (2006) 1115-1124.

- 771 [66] V.A. Payne, W.S. Au, S.L. Gray, E.D. Nora, S.M. Rahman, R. Sanders, D.  
772 Hadaschik, J.E. Friedman, S. O'Rahilly, J.J. Rochford, Sequential regulation of  
773 diacylglycerol acyltransferase 2 expression by CAAT/enhancer-binding protein  
774 beta (C/EBP $\beta$ ) and C/EBP $\beta$  during adipogenesis, *J Biol Chem* 282 (2007) 21005-  
775 21014.
- 776 [67] N. Arai, H. Masuzaki, T. Tanaka, T. Ishii, S. Yasue, N. Kobayashi, T. Tomita, M.  
777 Noguchi, T. Kusakabe, J. Fujikura, K. Ebihara, M. Hirata, K. Hosoda, T. Hayashi,  
778 H. Sawai, Y. Minokoshi, K. Nakao, Ceramide and adenosine 5'-monophosphate-  
779 activated protein kinase are two novel regulators of 11 $\beta$ -hydroxysteroid  
780 dehydrogenase type 1 expression and activity in cultured preadipocytes,  
781 *Endocrinology* 148 (2007) 5268-5277.
- 782 [68] S. Sai, C.L. Esteves, V. Kelly, Z. Michailidou, K. Anderson, A.P. Coll, Y.  
783 Nakagawa, T. Ohzeki, J.R. Seckl, K.E. Chapman, Glucocorticoid regulation of the  
784 promoter of 11 $\beta$ -hydroxysteroid dehydrogenase type 1 is indirect and requires  
785 C/EBP $\beta$ , *Mol Endocrinol* 22 (2008) 2049-2060.
- 786 [69] D. Ruffell, F. Mourkioti, A. Gambardella, P. Kirstetter, R.G. Lopez, N. Rosenthal,  
787 C. Nerlov, A CREB-C/EBP $\beta$  cascade induces M2 macrophage-specific gene  
788 expression and promotes muscle injury repair, *Proc Natl Acad Sci U S A* 106  
789 (2009) 17475-17480.
- 790 [70] B. Gorgoni, D. Maritano, P. Marthyn, M. Righi, V. Poli, C/EBP beta gene  
791 inactivation causes both impaired and enhanced gene expression and inverse  
792 regulation of IL-12 p40 and p35 mRNAs in macrophages, *J Immunol* 168 (2002)  
793 4055-4062.
- 794 [71] C.L. Esteves, V. Kelly, V. Begay, T.Y. Man, N.M. Morton, A. Leutz, J.R. Seckl,  
795 K.E. Chapman, Regulation of Adipocyte 11 $\beta$ -Hydroxysteroid Dehydrogenase  
796 Type 1 (11 $\beta$ -HSD1) by CCAAT/Enhancer-Binding Protein (C/EBP) beta Isoforms,  
797 *LIP and LAP, PLoS One* 7 (2012) e37953.
- 798 [72] J.J. Smink, V. Begay, T. Schoenmaker, E. Sterneck, T.J. de Vries, A. Leutz,  
799 Transcription factor C/EBPbeta isoform ratio regulates osteoclastogenesis through  
800 MafB, *EMBO J* 28 (2009) 1769-1781.
- 801 [73] J.J. Smink, A. Leutz, Rapamycin and the transcription factor C/EBPbeta as a  
802 switch in osteoclast differentiation: implications for lytic bone diseases, *J Mol Med*  
803 88 (2009) 227-233.
- 804 [74] M.J. Rauh, V. Ho, C. Pereira, A. Sham, L.M. Sly, V. Lam, L. Huxham, A.I.  
805 Minchinton, A. Mui, G. Krystal, SHIP represses the generation of alternatively  
806 activated macrophages, *Immunity* 23 (2005) 361-374.
- 807 [75] A. Arranz, C. Doxaki, E. Vergadi, Y. Martinez de la Torre, K. Vaporidi, E.D.  
808 Lagoudaki, E. Ieronymaki, A. Androulidaki, M. Venihaki, A.N. Margioris, E.N.  
809 Stathopoulos, P.N. Tsichlis, C. Tsatsanis, Akt1 and Akt2 protein kinases  
810 differentially contribute to macrophage polarization, *Proc Natl Acad Sci U S A* 109  
811 (2012) 9517-9522.
- 812 [76] A.G. Atanasov, L.G. Nashev, R.A. Schweizer, C. Frick, A. Odermatt, Hexose-6-  
813 phosphate dehydrogenase determines the reaction direction of 11 $\beta$ -hydroxysteroid  
814 dehydrogenase type 1 as an oxoreductase, *FEBS Lett* 571 (2004) 129-133.
- 815 [77] G. Banhegyi, A. Benedetti, R. Fulceri, S. Senesi, Cooperativity between 11 $\beta$ -  
816 hydroxysteroid dehydrogenase type 1 and hexose-6-phosphate dehydrogenase in  
817 the lumen of the endoplasmic reticulum, *J Biol Chem* 279 (2004) 27017-27021.
- 818 [78] I.J. Bujalska, N. Draper, Z. Michailidou, J.W. Tomlinson, P.C. White, K.E.  
819 Chapman, E.A. Walker, P.M. Stewart, Hexose-6-phosphate dehydrogenase confers

- 820 oxo-reductase activity upon 11 $\beta$ -hydroxysteroid dehydrogenase type 1, *J Mol*  
 821 *Endocrinol* 34 (2005) 675-684.
- 822 [79] E.A. Walker, A. Ahmed, G.G. Lavery, J.W. Tomlinson, S.Y. Kim, M.S. Cooper,  
 823 J.P. Ride, B.A. Hughes, C.H. Shackleton, P. McKiernan, E. Elias, J.Y. Chou, P.M.  
 824 Stewart, 11 $\beta$ -Hydroxysteroid Dehydrogenase Type 1 Regulation by Intracellular  
 825 Glucose 6-Phosphate Provides Evidence for a Novel Link between Glucose  
 826 Metabolism and Hypothalamo-Pituitary-Adrenal Axis Function, *J Biol Chem* 282  
 827 (2007) 27030-27036.
- 828 [80] R. Bertini, M. Bianchi, P. Ghezzi, Adrenalectomy sensitizes mice to the lethal  
 829 effects of Interleukin-1 and Tumor Necrosis Factor, *J Exp Med* 167 (1988) 1708-  
 830 1712.
- 831 [81] T.Y. Zhang, R.A. Daynes, Macrophages from 11 $\beta$ -hydroxysteroid dehydrogenase  
 832 type 1-deficient mice exhibit an increased sensitivity to lipopolysaccharide  
 833 stimulation due to TGF- $\beta$ -mediated up-regulation of SHIP1 expression, *J Immunol*  
 834 179 (2007) 6325-6335.
- 835 [82] S.J. McSweeney, P.W. Hadoke, A.M. Kozak, G.R. Small, H. Khaled, B.R. Walker,  
 836 G.A. Gray, Improved heart function follows enhanced inflammatory cell  
 837 recruitment and angiogenesis in 11 $\beta$ -HSD1-deficient mice post-MI, *Cardiovasc*  
 838 *Res* 88 (2010) 159-167.
- 839 [83] A.E. Coutinho, M. Gray, D.G. Brownstein, D.M. Salter, D.A. Sawatzky, S. Clay,  
 840 J.S. Gilmour, J.R. Seckl, J.S. Savill, K.E. Chapman, 11 $\beta$ -Hydroxysteroid  
 841 Dehydrogenase Type 1, But Not Type 2, Deficiency Worsens Acute Inflammation  
 842 and Experimental Arthritis in Mice, *Endocrinology* 153 (2012) 234-240.
- 843 [84] T. Kardon, S. Senesi, P. Marcolongo, B. Legeza, G. Banhegyi, J. Mandl, R.  
 844 Fulceri, A. Benedetti, Maintenance of luminal NADPH in the endoplasmic  
 845 reticulum promotes the survival of human neutrophil granulocytes, *FEBS Lett* 582  
 846 (2008) 1809-1815.
- 847 [85] G.R. Small, P.W. Hadoke, I. Sharif, A.R. Dover, D. Armour, C.J. Kenyon, G.A.  
 848 Gray, B.R. Walker, Preventing local regeneration of glucocorticoids by 11 $\beta$ -  
 849 hydroxysteroid dehydrogenase type 1 enhances angiogenesis, *Proc Natl Acad Sci*  
 850 *U S A* 102 (2005) 12165-12170.
- 851 [86] M. Wamil, J.H. Battle, S. Turban, T. Kipari, D. Seguret, R. de Sousa Peixoto, Y.B.  
 852 Nelson, D. Nowakowska, D. Ferenbach, L. Ramage, K.E. Chapman, J. Hughes,  
 853 D.R. Dunbar, J.R. Seckl, N.M. Morton, Novel fat depot-specific mechanisms  
 854 underlie resistance to visceral obesity and inflammation in 11 $\beta$ -hydroxysteroid  
 855 dehydrogenase type 1-deficient mice, *Diabetes* 60 (2011) 1158-1167.
- 856 [87] Z. Michailidou, S. Turban, E. Miller, X. Zou, J. Schrader, P.J. Ratcliffe, P.W.  
 857 Hadoke, B.R. Walker, J.P. Iredale, N.M. Morton, J.R. Seckl, Increased  
 858 Angiogenesis Protects against Adipose Hypoxia and Fibrosis in Metabolic  
 859 Disease-resistant 11 $\beta$ -Hydroxysteroid Dehydrogenase Type 1 (HSD1)-deficient  
 860 Mice, *J Biol Chem* 287 (2012) 4188-4197.
- 861 [88] R. Carter, J.M. Paterson, U. Tworowska, D.J. Stenvers, J.J. Mullins, J.R. Seckl,  
 862 M.C. Holmes, Hypothalamic-pituitary-adrenal axis abnormalities in response to  
 863 deletion of 11 $\beta$ -HSD1 is strain-dependent, *J Neuroendocrinol* 21 (2009) 879-  
 864 877.
- 865 [89] A. Sica, A. Mantovani, Macrophage plasticity and polarization: in vivo veritas, *J*  
 866 *Clin Invest* 122 (2012) 787-795.

- 867 [90] L.M. Sly, M.J. Rauh, J. Kalesnikoff, C.H. Song, G. Krystal, LPS-induced  
868 upregulation of SHIP is essential for endotoxin tolerance, *Immunity* 21 (2004) 227-  
869 239.
- 870 [91] N.M. Morton, J.M. Paterson, H. Masuzaki, M.C. Holmes, B. Staels, C. Fievet, B.R.  
871 Walker, J.S. Flier, J.J. Mullins, J.R. Seckl, Novel adipose tissue-mediated  
872 resistance to diet-induced visceral obesity in 11 $\beta$ -hydroxysteroid dehydrogenase  
873 type 1-deficient mice, *Diabetes* 53 (2004) 931-938.
- 874 [92] J.R. Seckl, N.M. Morton, K.E. Chapman, B.R. Walker, Glucocorticoids and 11 $\beta$ -  
875 hydroxysteroid dehydrogenase in adipose tissue, *Recent Prog Horm Res* 59 (2004)  
876 359-393.
- 877 [93] C. Edwards, Sixty years after Hench--corticosteroids and chronic inflammatory  
878 disease, *J Clin Endocrinol Metab* 97 (2012) 1443-1451.
- 879 [94] Y. Ichikawa, K. Yoshida, M. Kawagoe, E. Saito, Y. Abe, K. Arikawa, M. Homma,  
880 Altered equilibrium between cortisol and cortisone in plasma in thyroid  
881 dysfunction and inflammatory diseases, *Metabolism* 26 (1977) 989-997.
- 882 [95] Y. Kotelevtsev, M.C. Holmes, A. Burchell, P.M. Houston, D. Schmolli, P.  
883 Jamieson, R. Best, R. Brown, C.R.W. Edwards, J.R. Seckl, J.J. Mullins, 11 $\beta$ -  
884 hydroxysteroid dehydrogenase type 1 knockout mice show attenuated  
885 glucocorticoid inducible responses and resist hyperglycaemia on obesity or stress.,  
886 *Proc Natl Acad Sci USA* 94 (1997) 14924-14929.
- 887 [96] N.M. Morton, M.C. Holmes, C. Fiévet, B. Staels, A. Tailleux, J.J. Mullins, J.R.  
888 Seckl, Improved lipid and lipoprotein profile, hepatic insulin sensitivity, and  
889 glucose tolerance in 11 $\beta$ -hydroxysteroid dehydrogenase type 1 null mice, *J Biol*  
890 *Chem* 276 (2001) 41293-41300.
- 891 [97] P. Alberts, L. Engblom, N. Edling, M. Forsgren, G. Klingstrom, C. Larsson, Y.  
892 Ronquist-Nii, B. Ohman, L. Abrahmsen, Selective inhibition of 11 $\beta$ -  
893 hydroxysteroid dehydrogenase type 1 decreases blood glucose concentrations in  
894 hyperglycaemic mice, *Diabetologia* 45 (2002) 1528-1532.
- 895 [98] P. Alberts, C. Nilsson, G. Selen, L.O. Engblom, N.H. Edling, S. Norling, G.  
896 Klingstrom, C. Larsson, M. Forsgren, M. Ashkzari, C.E. Nilsson, M. Fiedler, E.  
897 Bergqvist, B. Ohman, E. Bjorkstrand, L.B. Abrahmsen, Selective inhibition of  
898 11 $\beta$ -hydroxysteroid dehydrogenase type 1 improves hepatic insulin sensitivity in  
899 hyperglycemic mice strains, *Endocrinology* 144 (2003) 4755-4762.
- 900 [99] M. Berthiaume, M. Laplante, W. Festuccia, Y. Gelinas, S. Poulin, J. Lalonde, D.R.  
901 Joannis, R. Thieringer, Y. Deshaies, Depot-specific modulation of rat  
902 intraabdominal adipose tissue lipid metabolism by pharmacological inhibition of  
903 11 $\beta$ -hydroxysteroid dehydrogenase type 1, *Endocrinology* 148 (2007) 2391-2397.
- 904 [100] M. Berthiaume, M. Laplante, W.T. Festuccia, K. Cianflone, L.P. Turcotte, D.R.  
905 Joannis, G. Olivecrona, R. Thieringer, Y. Deshaies, 11 $\beta$ -HSD1 inhibition  
906 improves triglyceridemia through reduced liver VLDL secretion and partitions  
907 lipids toward oxidative tissues, *Am J Physiol Endocrinol Metab* 293 (2007) E1045-  
908 1052.
- 909 [101] M. Berthiaume, M. Laplante, W.T. Festuccia, J.P. Berger, R. Thieringer, Y.  
910 Deshaies, Preliminary report: pharmacologic 11 $\beta$ -hydroxysteroid dehydrogenase  
911 type 1 inhibition increases hepatic fat oxidation in vivo and expression of related  
912 genes in rats fed an obesogenic diet, *Metabolism* 59 (2010) 114-117.
- 913 [102] J. Rosenstock, S. Banarar, V.A. Fonseca, S.E. Inzucchi, W. Sun, W. Yao, G.  
914 Hollis, R. Flores, R. Levy, W.V. Williams, J.R. Seckl, R. Huber, The 11 $\beta$ -  
915 hydroxysteroid dehydrogenase type 1 inhibitor INCB13739 improves

- hyperglycemia in patients with type 2 diabetes inadequately controlled by metformin monotherapy, *Diabetes Care* 33 (2010) 1516-1522.
- [103] P.U. Feig, S. Shah, A. Hermanowski Vosatka, D. Plotkin, M.S. Springer, S. Donahue, C. Thach, E.J. Klein, E. Lai, K.D. Kaufman, Effects of an 11 $\beta$ -hydroxysteroid dehydrogenase type 1 inhibitor, MK-0916, in patients with type 2 diabetes mellitus and metabolic syndrome, *Diabetes Obes Metab* 13 (2011) 498-504.
- [104] S. Shah, A. Hermanowski-Vosatka, K. Gibson, R.A. Ruck, G. Jia, J. Zhang, P.M.T. Hwang, N.W. Ryan, R.B. Langdon, P.U. Feig, Efficacy and safety of the selective 11 $\beta$ -HSD-1 inhibitors MK-0736 and MK-0916 in overweight and obese patients with hypertension, *Journal of the American Society of Hypertension* 5 (2011) 166-176.
- [105] M.M. Veniant, C. Hale, R. Komorowski, M.M. Chen, D.J. St Jean, C. Fotsch, M. Wang, Time of the day for 11 $\beta$ -HSD1 inhibition plays a role in improving glucose homeostasis in DIO mice, *Diabetes Obes Metab* 11 (2009) 109-117.
- [106] C.S. Wyrwoll, M.C. Holmes, J.R. Seckl, 11 $\beta$ -Hydroxysteroid dehydrogenases and the brain: From zero to hero, a decade of progress, *Frontiers in Neuroendocrinology* 32 (2011) 265-286.
- [107] J. Buren, S.-A. Bergstrom, E. Loh, I. Soderstrom, T. Olsson, C. Mattsson, Hippocampal 11 $\beta$ -hydroxysteroid dehydrogenase type 1 mRNA expression has a diurnal variability which is lost in the obese Zucker rat, *Endocrinology* 148 (2007) 2716-2722.
- [108] H.J. Harris, Y. Kotelevtsev, J.J. Mullins, J.R. Seckl, M.C. Holmes, Intracellular regeneration of glucocorticoids by 11 $\beta$ -hydroxysteroid dehydrogenase (11 $\beta$ -HSD)-1 plays a key role in regulation of the hypothalamic-pituitary-adrenal axis: analysis of 11 $\beta$ -HSD-1 deficient mice, *Endocrinology* 142 (2001) 114-120.
- [109] P.H. Bisschop, M.J. Dekker, W. Osterthun, J. Kwakkel, J.J. Anink, A. Boelen, U.A. Unmehopa, J.W. Koper, S.W. Lamberts, P.M. Stewart, D.F. Swaab, E. Fliers, Expression of 11 $\beta$ -hydroxysteroid dehydrogenase type 1 in the human hypothalamus, *J Neuroendocrinol* (2013) Jan 3. doi: 10.1111/jne.12017. [Epub ahead of print].
- [110] G.G. Lavery, A.E. Zielinska, L.L. Gathercole, B. Hughes, N. Semjonous, P. Guest, K. Saqib, M. Sherlock, G. Reynolds, S.A. Morgan, J.W. Tomlinson, E.A. Walker, E.H. Rabbitt, P.M. Stewart, Lack of Significant Metabolic Abnormalities in Mice with Liver-Specific Disruption of 11 $\beta$ -Hydroxysteroid Dehydrogenase Type 1, *Endocrinology* 153 (2012) 3236-3248.
- [111] Z. Michailidou, M.D. Jensen, D.A. Dumesic, K.E. Chapman, J.R. Seckl, B.R. Walker, N.M. Morton, Omental 11 $\beta$ -hydroxysteroid dehydrogenase 1 correlates with fat cell size independently of obesity, *Obesity* 15 (2007) 1155-1163.
- [112] M.J. Lee, S.K. Fried, S.S. Mundt, Y. Wang, S. Sullivan, A. Stefanni, B.L. Daugherty, A. Hermanowski-Vosatka, Depot-specific regulation of the conversion of cortisone to cortisol in human adipose tissue, *Obesity (Silver Spring)* 16 (2008) 1178-1185.
- [113] S. Cinti, G. Mitchell, G. Barbatelli, I. Murano, E. Ceresi, E. Faloia, S. Wang, M. Fortier, A.S. Greenberg, M.S. Obin, Adipocyte death defines macrophage localization and function in adipose tissue of obese mice and humans, *J Lipid Res* 46 (2005) 2347-2355.
- [114] M. Jernas, J. Palming, K. Sjoholm, E. Jennische, P.A. Svensson, B.G. Gabrielsson, M. Levin, A. Sjogren, M. Rudemo, T.C. Lystig, B. Carlsson, L.M.

- 965 Carlsson, M. Lonn, Separation of human adipocytes by size: hypertrophic fat cells  
 966 display distinct gene expression, *FASEB J* 20 (2006) 1540-1542.
- 967 [115] H. Masuzaki, J. Paterson, H. Shinyama, N.M. Morton, J.J. Mullins, J.R. Seckl,  
 968 J.S. Flier, A transgenic model of visceral obesity and the metabolic syndrome,  
 969 *Science* 294 (2001) 2166-2170.
- 970 [116] H. Masuzaki, H. Yamamoto, C.J. Kenyon, J.K. Elmquist, N.M. Morton, J.M.  
 971 Paterson, H. Shinyama, M.G. Sharp, S. Fleming, J.J. Mullins, J.R. Seckl, J.S. Flier,  
 972 Transgenic amplification of glucocorticoid action in adipose tissue causes high  
 973 blood pressure in mice, *J Clin Invest* 112 (2003) 83-90.
- 974 [117] E.E. Kershaw, N.M. Morton, H. Dhillon, L. Ramage, J.R. Seckl, J.S. Flier,  
 975 Adipocyte-specific glucocorticoid inactivation protects against diet-induced  
 976 obesity, *Diabetes* 54 (2005) 1023-1031.
- 977 [118] S.A. Hawley, M.D. Fullerton, F.A. Ross, J.D. Schertzer, C. Chevtzoff, K.J.  
 978 Walker, M.W. Pegg, D. Zibrova, K.A. Green, K.J. Mustard, B.E. Kemp, K.  
 979 Sakamoto, G.R. Steinberg, D.G. Hardie, The Ancient Drug Salicylate Directly  
 980 Activates AMP-Activated Protein Kinase, *Science* 336 (2012) 918-922.
- 981 [119] N. Hosogai, A. Fukuhara, K. Oshima, Y. Miyata, S. Tanaka, K. Segawa, S.  
 982 Furukawa, Y. Tochino, R. Komuro, M. Matsuda, I. Shimomura, Adipose tissue  
 983 hypoxia in obesity and its impact on adipocytokine dysregulation, *Diabetes* 56  
 984 (2007) 901-911.
- 985 [120] I. Stuart Wood, F.t.P.r. de Heredia, B. Wang, P. Trayhurn, Cellular hypoxia and  
 986 adipose tissue dysfunction in obesity, *Proceedings of the Nutrition Society* 68  
 987 (2009) 370.
- 988 [121] N. Halberg, T. Khan, M.E. Trujillo, I. Wernstedt-Asterholm, A.D. Attie, S.  
 989 Sherwani, Z.V. Wang, S. Landskroner-Eiger, S. Dineen, U.J. Magalang, R.A.  
 990 Brekken, P.E. Scherer, Hypoxia-inducible factor 1 $\alpha$  induces fibrosis and  
 991 insulin resistance in white adipose tissue, *Mol Cell Biol* 29 (2009) 4467-4483.
- 992 [122] A. Hermanowski-Vosatka, J.M. Balkovec, K. Cheng, H.Y. Chen, M. Hernandez,  
 993 G.C. Koo, C.B. Le Grand, Z. Li, J.M. Metzger, S.S. Mundt, H. Noonan, C.N.  
 994 Nunes, S.H. Olson, B. Pikounis, N. Ren, N. Robertson, J.M. Schaeffer, K. Shah,  
 995 M.S. Springer, A.M. Strack, M. Strowski, K. Wu, T. Wu, J. Xiao, B.B. Zhang,  
 996 S.D. Wright, R. Thieringer, 11 $\beta$ -HSD1 inhibition ameliorates metabolic syndrome  
 997 and prevents progression of atherosclerosis in mice, *J Exp Med* 202 (2005) 517-  
 998 527.
- 999 [123] M.J. Luo, R. Thieringer, M.S. Springer, S.D. Wright, A. Hermanowski-Vosatka,  
 1000 A. Plump, J.M. Balkovec, K. Cheng, G.J. Ding, D.W. Kawka, G.C. Koo, C.B. Le  
 1001 Grand, Q. Luo, M.M. Maletic, L. Malkowitz, K. Shah, I. Singer, S.T. Waddell,  
 1002 K.K. Wu, J. Yuan, J. Zhu, S. Stepaniants, X. Yang, P.Y. Lum, I.M. Wang, 11 $\beta$ -  
 1003 HSD1 inhibition reduces atherosclerosis in mice by altering pro-inflammatory gene  
 1004 expression in the vasculature, *Physiol Genomics* 45 (2013) 47-57.
- 1005 [124] T.M.J. Kipari, T.-Y. Man, P.W.F. Hadoke, J.S. Savill, K.E. Chapman, J.R. Seckl,  
 1006 11 $\beta$ -hydroxysteroid dehydrogenase type 1 deficiency in bone marrow-derived cells  
 1007 reduces atherosclerosis, *FASEB J* in press (2013).
- 1008 [125] G.A. Deuchar, D. McLean, P.W.F. Hadoke, D.G. Brownstein, D.J. Webb, J.J.  
 1009 Mullins, K. Chapman, J.R. Seckl, Y.V. Kotelevtsev, 11 $\beta$ -Hydroxysteroid  
 1010 Dehydrogenase Type 2 Deficiency Accelerates Atherogenesis and Causes  
 1011 Proinflammatory Changes in the Endothelium in Apoe $^{-/-}$  Mice, *Endocrinology* 152  
 1012 (2011) 236-246.

- 1013 [126] R.S. Hardy, E.H. Rabbitt, A. Filer, P. Emery, M. Hewison, P.M. Stewart, N.  
1014 Gittoes, C.D. Buckley, K. Raza, M.S. Cooper, Local and systemic glucocorticoid  
1015 metabolism in inflammatory arthritis, *Ann Rheum Dis* 67 (2008) 1204-1210.
- 1016 [127] M. Schmidt, C. Weidler, H. Naumann, S. Anders, J. Scholmerich, R.H. Straub,  
1017 Reduced capacity for the reactivation of glucocorticoids in rheumatoid arthritis  
1018 synovial cells: Possible role of the sympathetic nervous system?, *Arthritis Rheum*  
1019 52 (2005) 1711-1720.
- 1020 [128] R.S. Hardy, A. Filer, M.S. Cooper, G. Parsonage, K. Raza, D.L. Hardie, E.H.  
1021 Rabbitt, P.M. Stewart, C.D. Buckley, M. Hewison, Differential expression,  
1022 function and response to inflammatory stimuli of 11 $\beta$ -hydroxysteroid  
1023 dehydrogenase type 1 in human fibroblasts: a mechanism for tissue-specific  
1024 regulation of inflammation, *Arthritis Res Ther* 8 (2006) R108.
- 1025 [129] N. Olsen, T. Sokka, C.L. Seehorn, B. Kraft, K. Maas, J. Moore, T.M. Aune, A  
1026 gene expression signature for recent onset rheumatoid arthritis in peripheral blood  
1027 mononuclear cells, *Ann Rheum Dis* 63 (2004) 1387-1392.
- 1028 [130] C.S. Haas, C.J. Creighton, X. Pi, I. Maine, A.E. Koch, G.K. Haines, S. Ling,  
1029 A.M. Chinnaiyan, J. Holoshitz, Identification of genes modulated in rheumatoid  
1030 arthritis using complementary DNA microarray analysis of lymphoblastoid B cell  
1031 lines from disease-discordant monozygotic twins, *Arthritis Rheum* 54 (2006) 2047-  
1032 2060.
- 1033 [131] S. Suzuki, H. Tsubochi, H. Ishibashi, T. Suzuki, T. Kondo, H. Sasano, Increased  
1034 expression of 11 $\beta$ -hydroxysteroid dehydrogenase type 2 in the lungs of patients  
1035 with acute respiratory distress syndrome, *Pathol Int* 53 (2003) 751-756.
- 1036 [132] M.J. Young, Mechanisms of mineralocorticoid receptor-mediated cardiac fibrosis  
1037 and vascular inflammation, *Curr Opin Nephrol Hypertens* 17 (2008) 174-180.
- 1038 [133] J.M. Luther, P. Luo, Z. Wang, S.E. Cohen, H.S. Kim, A.B. Fogo, N.J. Brown,  
1039 Aldosterone deficiency and mineralocorticoid receptor antagonism prevent  
1040 angiotensin II-induced cardiac, renal, and vascular injury, *Kidney Int* 82 (2012)  
1041 643-651.
- 1042 [134] E.R. de Kloet, J.M. Reul, W. Sutanto, Corticosteroids and the brain, *J Steroid*  
1043 *Biochem Mol Biol* 37 (1990) 387-394.
- 1044 [135] J.W. Funder, Mineralocorticoid receptor activation and oxidative stress,  
1045 *Hypertension* 50 (2007) 840-841.
- 1046 [136] C. Guo, V. Ricchiuti, B.Q. Lian, T.M. Yao, P. Coutinho, J.R. Romero, J. Li, G.H.  
1047 Williams, G.K. Adler, Mineralocorticoid receptor blockade reverses obesity-  
1048 related changes in expression of adiponectin, peroxisome proliferator-activated  
1049 receptor-gamma, and proinflammatory adipokines, *Circulation* 117 (2008) 2253-  
1050 2261.
- 1051 [137] R.A. Frieler, H. Meng, S.Z. Duan, S. Berger, G. Schutz, Y. He, G. Xi, M.M.  
1052 Wang, R.M. Mortensen, Myeloid-Specific Deletion of the Mineralocorticoid  
1053 Receptor Reduces Infarct Volume and Alters Inflammation During Cerebral  
1054 Ischemia, *Stroke* 42 (2011) 179-185.

# FIGURE LEGENDS

## **Figure 1. 11 $\beta$ -HSD1 is induced upon macrophage differentiation**

Expression of 11 $\beta$ -HSD1 is negligible in human monocytes, but is induced on differentiation into macrophages. Polarisation of macrophages to an M1 phenotype further induces 11 $\beta$ -HSD1 whereas polarisation to an M2 phenotype has no further effect on expression. Differentiation of monocytes into macrophages in the presence of IL-4 and/or IL-13 further induces 11 $\beta$ -HSD1 (see text for details).

1063

1064 ***Figure 2. Macrophage polarisation is associated with a switch in energy metabolism***

1065 M1 macrophages show a predominantly glycolytic metabolism. High levels of glucose-  
1066 6-phosphate (G6P) may ensure a ready supply of NADPH cofactor to 11! -HSD1,  
1067 driving high conversion of cortisone (E) to cortisol (F). M2 polarised macrophages are  
1068 oxidative, with lower levels of glycolysis and lower levels of 11! -HSD1 converting E  
1069 to F. Whether changes in energy metabolism drive changes in macrophage 11! -HSD1  
1070 expression is currently unknown (see text for details).

1071

1072 ***Figure 3. Effects of 11!-HSD1 deficiency/inhibition on acute and chronic***  
1073 ***inflammation***

1074 Deficiency or inhibition of 11! -HSD1 worsens or exacerbates acute inflammation, but  
1075 may also promote its successful resolution. During chronic metabolic inflammation  
1076 (obesity, atherosclerosis, diabetes), 11! -HSD1 deficiency/inhibition is beneficial,  
1077 reducing inflammatory cell recruitment to sites of inflammation and promoting insulin  
1078 sensitisation. However, during chronic non-resolving inflammation, the pro-angiogenic,  
1079 pro-fibrotic phenotype of 11! -HSD1 deficiency/inhibition may worsen tissue damage  
1080 (see text for details).

1081



Figure 1






Figure 2



M1 polarised macrophage


↑ Glycolysis  
 ↓ O<sub>2</sub> consumption  
 ↑ 11 $\beta$ -HSD1



M2 polarised macrophage

↓ Glycolysis  
 ↑ O<sub>2</sub> consumption  
 ↓ 11 $\beta$ -HSD1

Figure 3



## Highlights

- !! 11 $\beta$ -HSD1 converts inert glucocorticoids into active forms, amplifying glucocorticoid action
- !! 11 $\beta$ -HSD1 is markedly induced by pro-inflammatory cytokines
- !! 11 $\beta$ -HSD1 deficiency/inhibition worsens acute inflammation
- !! 11 $\beta$ -HSD1 inhibition reduces inflammation in obesity or atherosclerosis
- !! an increased angiogenic response may underlie some of the benefits